



## Water pollution influencing contamination of *Vibrio* bacteria in the coastal aquaculture area of Chanthaburi and Trat Provinces

Jakkapan Potipat<sup>1\*</sup>, Chawanrat Somnuek<sup>1</sup> and Sutthinee Mekprayoon<sup>2</sup>

<sup>1</sup>Department of Environmental Science, Faculty of Science and Technology, Rambhai Barni Rajabhat University, Chanthaburi 22000, THAILAND

<sup>2</sup>Department of Chemistry, Faculty of Science and Technology, Rambhai Barni Rajabhat University, Chanthaburi 22000, THAILAND

\*Corresponding author: jakkapan.p@rbru.ac.th

### ABSTRACT

Since 1982, the Eastern coastal area of Thailand has been developed from the Eastern Seaboard Project (ESP) to the Eastern Economic Corridor (EEC). The marine ecosystem of Chanthaburi and Trat Provinces was polluted by anthropogenic activities such as agriculture, transportation, tourism, fisheries, and urban communities. This study aims to investigate the marine environmental quality and the contamination of the *Vibrio* bacteria (*V. cholerae*, *V. parahaemolyticus* and *V. vulnificus*) in the coastal aquaculture area of Chanthaburi and Trat Provinces. Environmental sampling areas were designated at seven stations eastward from Tamai to Klonyai districts (about 150 km long stretch). The physicochemical parameters, including temperature, salinity, conductivity, pH, dissolved oxygen (DO), NH<sub>3</sub>, and major anions e.g. NO<sub>2</sub><sup>-</sup>, PO<sub>4</sub><sup>3-</sup> of seawater samples were measured at designated stations. Marine samples, including cockles (*Anadara granosa*), mussels (*Perna viridis*), oysters (*Saccostrea cucullata*) and white shrimp (*Litopenaeus vannamei*) were collected from aquaculture areas located in the estuarine ecosystem with simple random sampling. Our studies revealed that major inorganic substance concentrations followed this consequence order PO<sub>4</sub><sup>3-</sup> > NH<sub>3</sub> > NO<sub>2</sub><sup>-</sup>. The physicochemical properties indicated that seawater quality has been varied within the marine quality standard class III for aquaculture. The prevalence of *V. parahaemolyticus* and *V. vulnificus* showed in all bivalve samples were detected at 0.36 to 4.30 MPN/g and below the detection limit (0.3 MPN/g), respectively, whereas *V. cholerae* was not detected. This study also concluded that the periods of environmental sampling did not significantly influence the seawater quality and the level of *Vibrio* contamination. However, the difference of infection rates for *V. parahaemolyticus* and *V. vulnificus* depended on the marine species.

**Keywords:** Water pollution, *Vibrio* bacteria, Coastal aquaculture area

### INTRODUCTION

*Vibrio* is a Gram-negative bacteria, typically with lipopolysaccharide in the outer membrane, that belongs to the Proteobacteria phylum, Gammaproteobacteria class, the most diverse class of Gram-negative bacteria. The Vibrionaceae family comprises aquatic bacteria that mostly thrive in warm waters and tolerate various salinity levels, including freshwater, brackish, and marine waters [1].

Nowadays, the global aquaculture sector has grown continuously, and it is currently an important contributor to aquatic animal protein for human consumption [2]. The Thai government has to decide how they will control their national aquaculture production. Multiple bivalve species are economically important both natural and farmed populations, while white shrimp are produced mainly for export [3, 4]. Foodborne infections with the genus *Vibrio* are a serious

problem in Thailand and a major cause of gastroenteritis, particularly from traditional consumption of raw or undercooked seafood. Most of these patients are affected by *V. parahaemolyticus* and *V. cholerae* and to a lesser extent by *V. vulnificus* [5]. *Vibrios* belong to the microbiota of infectious bivalves, which can concentrate bacteria in their edible tissues and body fluids, including the hemolymph [6]. The aquacultural harvests from the Gulf of Thailand were more contaminated than *Vibrio*, while bivalves and white shrimps showed a high frequency of contamination [7-9].

The marine mollusks showed a great correlation to environmental variability investigation in the coastal areas and river mouth ecosystems [10]. Bivalve molluscs as bioindicator organisms have been used to assess pollution levels in the aquatic environment [11]. The bivalve species, including oysters (*Saccostrea*

*cucullata*), mussels (*Perna viridis*), and cockles (*Anadara granosa*) are abundance species and commercial seafood in the Eastern part of Thailand [7, 12]. Whereas the white shrimp (*Litopenaeus vannamei*) is used to focus on the potential *Vibrio* disease, due to *Vibrio* is recognized as a major cause of seafood-borne illness [13, 14].

Coastal water pollution is an increasingly significant environmental effect and public health illness [15]. Untreated wastewater and urban runoff are the principal sources of marine pollution [16]. These pollutants can inhibit the growth, reproduction, and survival of flora and fauna, leading to a decline in biodiversity. Physicochemical parameters are a limiting factor in the marine ecosystem and the most important environmental variable [17]. The various physicochemical measurements carried out in the coastal seawaters reveal the relationships between environmental quality and intensive anthropogenic activities [18, 19]. Hence, the physicochemical properties of seawater, including temperature, salinity, conductivity, pH, dissolved oxygen (DO), ammonia (NH<sub>3</sub>), and major anions (NO<sub>2</sub><sup>-</sup>, PO<sub>4</sub><sup>3-</sup>) are fundamental indicators for the assessment of changing the marine environment and reflecting land use development. Numerous indices regarding seawater quality and pollution status were conducted in Thailand, while the operation of these parameters was followed by a manual seawater sampling and analysis of PCD Thailand and international standards [20, 21].

The inner eastern coastal area of Thailand is the most important coastal aquaculture area and is a central source of recreation, as well as travel destinations, conservation areas, and comfort zones. In recent years, Thailand's Special Economic Zone (SEZ) was approved by the Thai government in 2015, covering the Chanthaburi and Trat coastal areas [22]. These marine zones have

seen significant spatial development in the past decade. As a result, they have transformed into important shrimp farming areas, tourism hotspots, and urban communities. Several reports indicated that anthropogenic activities as a source of water pollution are fed back into the marine ecosystem [23-26].

Here, we designed an analysis of water pollution influencing contamination and abundance of *Vibrio* bacteria by collecting marine samples along the Chanthaburi and Trat coastal areas from 2014 to 2022. Precisely, we determined the correlation between water quality parameters and the infection of *Vibrio* in commercial marine seafood with the statistical data.


## MATERIALS AND METHODS

### Environmental sampling station

The eastern coastal areas cover two provinces, namely Chanthaburi and Trat Provinces, which were considered the locations for the marine sampling (Figure 1). The seawater samples were selected to represent environmental quality at designated stations. All study sites were recorded by Global Positioning System (GPS) (SD ± 5 m) as shown in Table 1.

Three bivalve species, including cockles (*Anadara granosa*), mussels (*Perna viridis*), and oysters (*Saccostrea cucullata*), as well as white shrimps (*Litopenaeus vannamei*) were collected from coastal aquaculture areas using simple random sampling. The surface seawater samples were collected at a depth of 100 cm using a vertical sampling procedure for physicochemical analysis.

All species samples were dredged and cleaned with seawater and stored in 4.0 °C iceboxes to the medical laboratory within 24 hours after collection.



**Figure 1** The map of sampling stations along the coastal area of Chanthaburi and Trat Provinces.

**Table 1** Use details of the sampling stations.

Stations	Latitude	Longitude	Description
Chanthaburi coastal areas			
Tha Mai (A)	12.621421°	102.004851°	Tourism landmarks and fisheries
Mueang Chanthaburi (B)	12.606965°	102.104579°	Several anthropogenic activities (Urban community, transportation and oxidation pond treatment)
Laem Sing (C)	12.481633°	102.073803°	Aquaculture in river mouth areas
Khlung (D)	12.454718°	102.221408°	Agriculture, homestay and fisheries
Trat coastal areas			
Mueang Trat (E)	12.243642°	102.515117°	Urban community, disposal site and fisheries
Laem Ngop (F)	12.225371°	102.369298°	Eco-tourism and fisheries
Khlong Yai (G)	12.200052°	102.296142°	Border trade and fisheries

**Table 2** The vulnerability assessment of the coastal marine ecosystem of Chanthaburi and Trat Provinces.

Variables	Chanthaburi	Trat
Habitat		
In-shore shallow Seawater	Low	Low to Middle
Sandy beaches	Low	Middle
Rocky beaches	Low	Middle
Inter-tidal mudflats	Middle	Middle to High
Estuaries	Very high	Very high
Seagrass beds	Loss	Loss
Mangroves	Very high	Very high
Selected species		
Cetaceans (3 species)	Very high	Very high
Dugong	Loss	Loss
Migratory shorebirds	Low	Low
Red-backed sea eagle	Low to Middle	Low to Middle

Source: Modification from [34].

### Physicochemical measurement

The physicochemical properties were recorded in situ in each station using the electrochemical analyzer (Consort C 932) and salinometer which were calibrated before use. Dissolved Oxygen (DO) is measured by azide modification of the Winkler's titration [20]. Direct nesslerization was performed for NH<sub>3</sub> detection, whereas the colorimetric technique was used to determine nitrite (NO<sub>2</sub><sup>-</sup>) and phosphate (PO<sub>4</sub><sup>3-</sup>) [27, 28].

### Bacteriological analysis

Marine samples were prepared by rinsing and scrubbing them under running deionized water to remove debris from the shell, and they were opened using a sterile knife. The edible tissues in an equal amount of phosphate-buffered saline (PBS; pH 7.4) were homogenized in a sterile blender for 90 s [29].

The observation of *Vibrio* contamination was prepared following methods outlined in the Bacteriological Analytical Manual (BAM) for food sampling/preparation [30]. *V. cholerae*, *V. parahaemolyticus*, and *V. vulnificus*

were isolated from marine species by cultured in alkaline peptone water (APW) and thiosulfate citrate bile sucrose (TCBS), which was modified from BAM [31]. Each APW culture sample was sub-cultured on TCBS agar and incubated at 35°C for 18 - 24 hours to observe fermentative colonies. *Vibrio* densities in contaminated samples were calculated using the scored MPN table, and each dilution was measured in triplicate [32].

### Data analysis and quality control

A prevalence of *Vibrio* for each of the four species was tested for significant differences by analysis of Chi-square;  $p < 0.05$  was accepted for significant value. The influence of time on study periods and the physicochemical parameters of seawaters were analyzed using one-way analysis of variance (ANOVA) with  $p < 0.05$  was classified as statistically significant. Post hoc comparisons, LSD (Least Significant Difference) were applied to report differences among the sampling periods;  $p < 0.05$  was accepted as a significant value. The statistical inferences for hypothesis



tests were confirmed with a 5% significance level. All statistical analyses were performed using SPSS 18.0 software (Serial No.5083337). The quality control procedures were implemented through rigorous standardization by the PCD (Pollution Control Department), in accordance with seawater analysis guidelines.

## RESULTS AND DISCUSSION


### *Land use of coastal area of Chanthaburi and Trat Provinces*

The eastern coastal region of Thailand has been an important part of the country's economic development. The real gross regional product (GRP) of the eastern region increased significantly from around 500,000 THB in 1994 to 1.7 million THB in 2018 [33]. The terrestrial area of Chanthaburi and Trat Provinces is the most agricultural landscape. It covers more than 60 percent of the total land use. The agricultural areas are a mixture of many kinds of tropical fruits known for their tasteful flavor. The coastal areas of Chanthaburi and Trat Provinces are affected by four major river basins, including the Wang-Ta-Nord, Chanthaburi, Welu, and Trat river basins, as shown in Table 2. Land use activities from the river basins during heavy rainfall may also significantly affect the marine coastal area and the dynamic of *Vibrio* spp. abundance [35].

### *Physicochemical parameters*

The physicochemical parameters are used to determine environmental quality along the coastal

areas of Chanthaburi and Trat Provinces. Table 3 reported the annual average of physicochemical properties from 2014, 2018 and 2022. The variation in seawater quality was elucidated with a Box plot diagram based on the minimum, maximum, first, median, and third quartile (Figure 2). The seawater quality monitoring in study sites could be categorized as a marine water quality standard class III. Additionally, the investigation of seawater samples revealed that the environmental quality of study areas was unpolluted.



**Figure 2** Box plot of physicochemical parameters of seawater samples (A: Temperature; B: Salinity; C: Conductivity; D: pH; E: Dissolved Oxygen (DO); F: Ammonia-Nitrogen; G: Nitrite-Nitrogen and H: Phosphate-Phosphorus).

**Table 3** Seawater quality along the coastal areas of Chanthaburi and Trat Provinces, Thailand of 2014, 2018 and 2022.

Physicochemical parameters	Year 2014	Year 2018	Year 2022	Criterion*
Temperature (°C)	29 - 33	30 - 34	28 - 34	a
Salinity (g/L)	31.14 ± 1.57	32.14 ± 1.35	30.86 ± 2.19	b
Conductivity (µS/cm)	8 - 13	9 - 14	9 - 16	-
pH	10.29 ± 1.70	11.71 ± 1.98	12.71 ± 2.50	-
Dissolved Oxygen (mg/L)	18.39 - 32.45	23.54 - 40.26	19.55 - 40.11	7.0 - 8.5
Ammonia-Nitrogen (µg/L)	24.83 ± 5.14	33.90 ± 5.44	32.10 ± 6.96	≥ 4
Nitrite-Nitrogen (µg/L)	7.04 - 8.11	7.09 - 8.95	7.36 - 8.09	≤ 70
Phosphate-Phosphorus (µg/L)	7.43 ± 0.37	7.66 ± 0.65	7.75 ± 0.34	55 <sup>c</sup>
	6.24 ± 0.65	6.72 ± 0.55	6.55 ± 0.44	≤ 45
	5.83 - 10.18	6.78 - 20.12	9.17 - 13.47	
	8.05 ± 1.68	11.34 ± 4.28	11.29 ± 1.48	
	7.15 - 18.21	6.32 - 12.13	8.09 - 11.37	
	10.67 ± 4.44	8.87 ± 1.89	9.39 ± 1.23	
	10.46 - 25.32	9.64 - 20.23	8.31 - 19.56	
	15.61 ± 5.17	13.09 ± 3.90	13.91 ± 4.39	

\*The announcement of the National Environment Board No.27 (B.E. 2549) regarding the specification of the standard of seawater

a - Naturally but changing by no more than 1 °C

b - Changing no more than 10 percent of the lowest value


c - AMEQC (ASEAN Marine Environmental Quality Criteria)

The statistical comparison did not reveal a significant difference between the seawater quality and the sampling periods ( $p > 0.05$ ), except for conductivity. Multiple comparisons (LSD) revealed a significant difference in conductivity values across the study periods period ( $p < 0.05$ ). The conductivity variation in seawater is affected by the presence of inorganic dissolved solids such as chloride, nitrate, phosphate, and other anions. Mostly, essential nutrients enter the marine environment through urban rinsing, irrigation drainage, and agricultural runoff, while the high nutrient levels in seawater may lead to phytoplankton abundance and eutrophication [36]. In this study, the sampling sites revealed the degree of variation in terrestrial water pollution, which is frequently observed in estuarine coastal areas due to locally accumulated contaminants from domestic wastewater, fertilizer usage in orchards, and natural watershed erosion. A septic tank with an anaerobic filter was recommended for practice in solving water pollution problems in urban areas [37]. Conversely,

nanomaterials, biosolids, and electrobiological techniques have efficiently removed nutrients from agricultural wastewater [38, 39].

*Contamination of Vibrio bacteria*

The contamination of *Vibrio* spp. was investigated in the marine samples collected along the coastal aquaculture area of Chanthaburi and Trat Provinces, which produced seafood products for Thailand's domestic consumption and also exported. The results showed the infection of *V. parahaemolyticus* in all marine species, but *V. cholerae* was not detected, except in three bivalve species that were contaminated with *V. vulnificus* (Table 4; Figure 3). However, abundance of *Vibrio* species was below the seafood safety standard for fishery products, indicating no potential risks for human consumption. The main contaminated source of *Vibrio* bacteria is the discharge from untreated wastewater, urban sewage and infectious leachate, especially during extremely heavy rainfall, which are very threatening to the downstream ecosystem [19].



**Figure 3** Distribution of *Vibrio* contamination in marine samples along the coastal areas of Chanthaburi and Trat Provinces from 2014 to 2022.

**Table 4** Contamination of *Vibrio* spp. (MPN/g) in 25 g samples of marine species along the coastal aquaculture areas of Chanthaburi and Trat Provinces (7 stations) from 2014 to 2022 ( $n = 63$ ).

Species	<i>V. cholerae</i>		<i>V. parahaemolyticus</i>		<i>V. vulnificus</i>	
	Prevalence (percent)	MPN/g	Prevalence (percent)	MPN/g	Prevalence (percent)	MPN/g
Cockles	-	n.d.	55.56	0.36 - 1.25	25.40	< 0.3
Mussels	-	n.d.	63.49	0.74 - 2.80	15.87	< 0.3
Oysters	-	n.d.	73.02	0.36 - 4.30	36.51	< 0.3
White shrimps	-	n.d.	47.62	< 0.3 - 3.0	-	n.d.
Safety levels*		n.d.		< 30/g (MPN)		< 30/g (MPN)

n.d. - not detected

\*FDA and EPA safety levels of fishery products in regulation and guidance.

**Table 5** Growth characteristics of *V. parahaemolyticus* in the marine environment

Variables	Optimum	Range
Temperature (°C)	37	5 - 43
pH	7.8 - 8.6	4.8 - 11
NaCl (%)	1.5 - 3.0	0.5 - 10
Water Activity (WA)	0.981	0.940 - 0.996
Atmosphere (O <sub>2</sub> consumption)	Aerobic	Aerobic-Anaerobic

Source: Modification from International Commission for Microbiological Specifications for Foods

The study found a high abundance of *V. parahaemolyticus* in all marine samples. The chi-square test showed a significant difference in the prevalence of *V. parahaemolyticus* in the marine species ( $p < 0.05$ ), and the sampling periods were not a significant difference ( $p > 0.05$ ). *V. parahaemolyticus* is commonly found in coastal areas and estuarine ecosystems, and it has been isolated in various marine animals [40]. *V. parahaemolyticus* is a mildly halophilic, mesophilic microorganism, and its general growth characteristics are shown in Table 5. Hence, the physicochemical variation can play an important role in the significant contamination of *V. parahaemolyticus* in marine habitats [41-44].

The contamination of *V. vulnificus* was detected in all samples of bivalves at below detection limit (0.3 MPN/g) but was not found in the white shrimp samples during the study periods (Table 4). The statistical analysis revealed a notable disparity in the infection levels of *V. vulnificus* in bivalve samples, with a significance level of  $p < 0.05$ . A microbiological survey concerned the levels of pathogenic *V. vulnificus* in marine bivalves across various global locations. The reporting indicated infection rates of 17.2% in oysters and 8.0% in mussels [45, 46]. The prevalence rate of *V. vulnificus* contamination is commonly observed in shellfish maturity within coastal areas, but in this study, *V. vulnificus* was not detected in the white shrimp samples. The confirmation of *V. vulnificus* appeared to have the same disease symptoms as naturally infected shrimp. However, the circulating hemocytes of *Litopenaeus*

*vannamei* play a crucial role in self-purification and activate several *Vibrio* pathogens [47]. Furthermore, physicochemical parameters, such as salinity and temperature, influence the occurrence of *V. vulnificus*, while the natural depuration pond was prepared to decrease *Vibrio* contamination [48, 49].

## CONCLUSIONS

The present study concludes that physicochemical parameters were within the seawater permissible standard at all sampling stations in the coastal aquaculture area of Chanthaburi and Trat Provinces. The major land use concerns that have polluted the marine coastal area from anthropogenic activities such as recreational renovation, aquaculture, and infrastructure improvement. *V. parahaemolyticus* is an abundant species in the aquaculture area and is commonly isolated from edible bivalves and white shrimp tissues. However, *V. parahaemolyticus* densities also included the permissible limit. The depuration processes were recommended to decontaminate the cultivated fishery products.

## ACKNOWLEDGEMENT

This research was financially supported by the national budget of the National Research Council of Thailand (NRCT) from 2012 to 2018. The authors also would like to greatly thank the Faculty of Science and Technology, Rambhai Barni Rajabhat University for the experimental facilities provided during the study period.

### Ethics approvals

Ethical approval was exempted from this study as it operated within the local fishing communities' methods.

### REFERENCES

1. Wright AC, Hill RT, Johnson JA, Roghman MC, Colwell RR, Morris JG. Distribution of *Vibrio vulnificus* in the Chesapeake Bay. *Appl Environ Microbiol.* 1996;62:717-24.
2. FAO (Food and Agriculture Organization of the United Nations). The state of world fisheries and aquaculture 2016. Rome: Food and Agriculture Organization of the United Nations; 2016.
3. Ferreira JG, Falconer L, Kittiwanch J, Ross L, Saurel C, Wellman K, et al. Analysis of production and environmental effects of Nile tilapia and white shrimp culture in Thailand. *Aquaculture.* 2015; 447:23-36.
4. Oliveira GFM, Couto MCM, Lima MF, Bomfim TCB. Mussels (*Perna perna*) as bioindicator of environmental contamination by *Cryptosporidium* species with zoonotic potential. *Int J Parasitol Parasites Wildl.* 2016;5:28-33.
5. FEHD (Food and Environmental Hygiene Department). *Vibrio* species in seafood. Hong Kong: The Government of the Hong Kong Special Administrative Region; 2005.
6. Destoumieux-Garzón D, Canesi L, Oyanedel D, Travers MA, Charrière GM, Pruzzo C, et al. *Vibrio*-bivalve interactions in health and disease. *Appl Microbiol Int.* 2020;22(10):4323-41.
7. Nakaguchi Y. Contamination by *Vibrio parahaemolyticus* and its virulent strains in seafood marketed in Thailand, Vietnam, Malaysia, and Indonesia. *Trop Med Health.* 2013;41(3):95-102.
8. Yokyingyong P. Risk assessment of *Vibrio parahaemolyticus* in white shrimp (*Litopenaeus vannamei*) from retail market [thesis]. Faculty of Veterinary Science: Chulalongkorn University; 2019.
9. Jearnsripong S, Chuanchuen R, Atwill ER. Assessment of bacterial accumulation and environmental factors in sentinel oysters and estuarine water quality from the Phang Nga estuary area in Thailand. *Int J Env Res Pub He.* 2018;15:1-17.
10. Nuangjui M, Pimpang B, Chulalaksananukul W, Glinwong C. Biomonitoring by using rapid-read pathogenic bacteria indicator in sediments and bivalve mollusks: southern Gulf of Thailand, a mangrove area case study. *Trends Sci.* 2023;20(4):1-14.
11. Boening DW. An evaluation of bivalves as biomonitors of heavy metals pollution in marine waters. *Environ Monit Assess.* 1999;55(3):459-70.
12. Cheevaporn V, Menasveta P. Water pollution and habitat degradation in the Gulf of Thailand. *Mar Pollut Bull.* 2003;47:43-51.
13. Gopal S, Otta SK, Kumar S, Karunasagar I, Nishibuchi M, Karunasagar I. The occurrence of *Vibrio* species in tropical shrimp culture environments; implications for food safety. *Int J Food Microbiol.* 2005;102(2):151-59.
14. Somboon M, Purivirojkul W, Limsuwan C, Chuchird N. Effect of *Vibrio* spp. In white feces infected shrimp in Chanthaburi, Thailand. *Fish Res Bull Kasetsart Univ.* 2012;36(1):7-15.
15. Pendergraft MA, Belda-Ferre P, Petras D, Morris CK, Mitts Ba, Aron AT, et al. Bacterial and chemical evidence of coastal water pollution from the Tijuana River in sea spray aerosol. *Environ Sci Technol.* 2023;57:4071-81.
16. Feddersen F, Boehm AB, Giddings SN, Wu X, Liden D. Modeling untreated wastewater evolution and swimmer illness for four wastewater infrastructure scenarios in the San Diego-Tijuana (US/MX) border region. *GeoHealth.* 2021;5:1-20.
17. Elahi N, Ahmed Q, Bat L, Yousuf F. Physicochemical parameters and seasonal variation of coastal water from Balochistan coast, Pakistan. *J Coast Life Med.* 2015;3(3):199-203.
18. Narendra Babu K, Omana PK, Mohan M. Water and sediment quality of Ashtamudi estuary, a Ramsar site, southwest coast of India-a statistical appraisal. *Environ Monit Assess.* 2010;165:307-19.
19. Bonisławska M, Nędzarek A, Rybczyk A, Tański A. Influence of anthropogenic pollution on the physicochemical conditions of the waters of the lower section of the Sapólna River. *Water.* 2024; 16(1):w16010035.
20. APHA (American Public Health Association). Standard methods for the examination of water and wastewater. 20th ed. Washington DC: APHA, AWWA & WEF; 1998.
21. PCD (Pollution Control Department). Manual seawater sampling and analysis. Bangkok: Teachers' Council Printing House; 2001.
22. Kanjananil B. The role of the state in the development of Special Economic Zone: the case study of Trat. *Rajapark J.* 2018;12(26):78-89.
23. Sarakhan N. The environment education model for sustainable mangrove forest management in the Eastern part of Thailand. *App Environ Res.* 2014;41(4):29-40.



24. PCD (Pollution Control Department). Thailand state of pollution report 2015. Bangkok: Ministry of Natural Resources and Environment; 2015.
25. PCD (Pollution Control Department). Thailand state of pollution report 2016. Bangkok: Ministry of Natural Resources and Environment; 2016.
26. Kunsook C, Dumrongrojwathana P. Species diversity and abundance of marine crabs (Portunidae: Decapoda) from a collapsible crab trap fishery at Kung Krabaen Bay, Chanthaburi Province, Thailand. *Trop Life Sci Res.* 2017;28(1):45-67.
27. Traichaiyaporn S. Water quality analysis. Chiang Mai: Faculty of Science, Chiang Mai University; 2000.
28. Daungsavat M, Somsiri J. Water properties and analysis for fisheries research. Bangkok: Department of Fisheries, Ministry of Agriculture and Cooperatives; n.d.
29. Brumfield KD, Chen AJ, Gangwar M, Usmani M, Hasan NA, Jutla AS, et al. Environmental factors influencing occurrence of *Vibrio parahaemolyticus* and *Vibrio vulnificus*. *Appl Environ Microbiol.* 2023;89(6):1-19.
30. Andrews WH, Hammack TS. Chapter 1: Food sampling and preparation of sample homogenate. In: Jinneman K, Chair Member. *Bacteriological Analytical Manual (BAM)*. FDA; 2019. p. 1-10.
31. Kaysner CA, DePaola A, Jones J. Chapter 9: *Vibrio*. In: Jinneman K, Chair Member. *Bacteriological Analytical Manual (BAM)*. FDA; 2004. p. 1-41.
32. Miwa N, Nishio T, Arita Y, Kawamori F, Masuda T, Akiyama M. Evaluation of MPN method combined with PCR procedure for detection and enumeration of *Vibrio parahaemolyticus* in seafood. *Food Hyg Safe Sci*;2003;44(6):289-93.
33. Tontisirin N, Anantsuksomsri S. Economic development policies and land use changes in Thailand: from the Eastern Seaboard to the Eastern Economic Corridor. *Sustainability.* 2021;13:6153.
34. Bezuijen MR, Morgan C, Mather, RJ. A rapid vulnerability assessment of coastal habitats and selected species to climate risks in Chathaburi and Trat (Thailand), Koh Kong and Kampot (Cambodia) and Kien Giang, Ben Tre, Soc Trang and Can Gio (Vietnam). Gland, Switzerland: IUCN (International Union for Conservation of Nature); 2011.
35. Cantet F, Hervio-Heath D, Caro A, Le Menec C, Monteil C, Quéméré C, et al. Quantification of *Vibrio parahaemolyticus*, *Vibrio vulnificus* and *Vibrio cholerae* in French Mediterranean coastal lagoons. *Res Microbiol.* 2013;164(8):867-74.
36. Akinde SB, Obire O. *In-situ* physico-chemical properties of the Deep Atlantic Ocean water column and their implications on heterotrophic bacterial distribution in the Gulf of Guinea. *Adv Appl Sci Res.* 2011;2(6):470-82.
37. Thongkao K. The assessment of domestic wastewater quality for the guideline of water treatment and remediation in urban area. *SNRU J Sci Technol.* 2016;8(1):155-62.
38. Janczukowicz W, Rodziewicz J. Water and wastewater management in agriculture. *Appl Sci.* 2024;14:2488.
39. Rusañescu CO, Rusañescu M, Constantin GA. Wastewater management in agriculture. *Water.* 2022;14:3351.
40. Yokyingyong P. Risk assessment of *Vibrio parahaemolyticus* in white shrimp (*Litopenaeus vannamei*) from retail market [thesis]. Faculty of Veterinary Science: Chulalongkorn University; 2019.
41. Blackwell KD, Oliver JD. Ecology of *Vibrio vulnificus*, *Vibrio cholerae* and *Vibrio parahaemolyticus* in North Carolina estuaries. *J Microbiol.* 2008;46(2):146-53.
42. Hsieh JL, Fries JS, Noble RT. Dynamics and predictive modeling of *Vibrio* spp. in the Neuse river estuary, North Carolina, USA. *Environ Microbiol.* 2008;10(1):57-64.
43. Chonsin K, Hemmanee B, Hongkul R, Changkaew K, Noisumdaeng P, Suthienkul O. Detection of bacteriophages specific to *Vibrio parahaemolyticus* in marine water samples, Surat Thani, Thailand. *Thai J Public Health.* 2021;5(1):7-15.
44. Padovan A, Siboni N, Kaestli M, King WL, Seymour JR, Gibb K. Occurrence and dynamics of potentially pathogenic vibrios in the wet-dry tropics of northern Australia. *Mar Environ Res.* 2021;169:105405.
45. Beneduce L, Vernile A, Spano G, Massa S, Lamacchia F, Oliver JD. Occurrence of *Vibrio vulnificus* in mussel farms from the Varano lagoon environment. *Lett Appl Microbiol.* 2010;51(4):443-49.
46. Kirs M, DePaola A, Fyfe R, Jones JL, Krantz J, Van Laanen A, et al. A survey of oysters (*Crassostrea gigas*) in New Zealand for *Vibrio parahaemolyticus* and *Vibrio vulnificus*. *Int J Food Microbiol.* 2011;147(2):149-53.
47. Liu CH, Yeh ST, Cheng SY, Chen JC. The immune response of the white shrimp *Litopenaeus vannamei* and its susceptibility to *Vibrio* infection in relation with the moult cycle. *Fish Shellfish Immun.* 2004;16(2):151-61.



48. Kelly MT. Effect of temperature and salinity on *Vibrio (Beneckea) vulnificus* occurrence in a Gulf Coast Environment. *Appl Environ Microbiol.* 1982; 44(4):820-24.
49. Chen YP. Reduction and management of *Vibrio vulnificus* in gulf coast oyster (*Crassostrea Virginica*) [dissertation]. School of Nutrition and Food Sciences: Louisiana State University; 1996.